
B-Cell-Specific Diversion of Glucose Carbon Utilization Reveals a Unique Vulnerability in B Cell Malignancies.

Journal: Cell

Publication Year: 2018

Authors: Gang Xiao, Lai N Chan, Lars Klemm, Daniel Braas, Zhengshan Chen, Huimin Geng, Qiuyi Chen Zhang, Ali Aghajanirofeh, Kadriye Nehir Cosgun, Teresa Sadras, Jaewoong Lee, Tamara Mirzapourzova, Ravi Salgia, Thomas Ernst, Andreas Hochhaus, Hassan Jumaa, Xiaoyan Jiang, David M Weinstock, Thomas G Graeber, Markus Muschen

PubMed link: 29551267

Funding Grants: Lgr5-mediated self-renewal in B cell selection and leukemia-initiation

Public Summary:

B cell activation during normal immune responses and oncogenic transformation impose increased metabolic demands on B cells and their ability to retain redox homeostasis. While the serine/threonine-protein phosphatase 2A (PP2A) was identified as a tumor suppressor in multiple types of cancer, our genetic studies revealed an essential role of PP2A in B cell tumors. Thereby, PP2A redirects glucose carbon utilization from glycolysis to the pentose phosphate pathway (PPP) to salvage oxidative stress. This unique vulnerability reflects constitutively low PPP activity in B cells and transcriptional repression of G6PD and other key PPP enzymes by the B cell transcription factors PAX5 and IKZF1. Reflecting B-cell-specific transcriptional PPP-repression, glucose carbon utilization in B cells is heavily skewed in favor of glycolysis resulting in lack of PPP-dependent antioxidant protection. These findings reveal a gatekeeper function of the PPP in a broad range of B cell malignancies that can be efficiently targeted by small molecule inhibition of PP2A and G6PD.

Scientific Abstract:

B cell activation during normal immune responses and oncogenic transformation impose increased metabolic demands on B cells and their ability to retain redox homeostasis. While the serine/threonine-protein phosphatase 2A (PP2A) was identified as a tumor suppressor in multiple types of cancer, our genetic studies revealed an essential role of PP2A in B cell tumors. Thereby, PP2A redirects glucose carbon utilization from glycolysis to the pentose phosphate pathway (PPP) to salvage oxidative stress. This unique vulnerability reflects constitutively low PPP activity in B cells and transcriptional repression of G6PD and other key PPP enzymes by the B cell transcription factors PAX5 and IKZF1. Reflecting B-cell-specific transcriptional PPP-repression, glucose carbon utilization in B cells is heavily skewed in favor of glycolysis resulting in lack of PPP-dependent antioxidant protection. These findings reveal a gatekeeper function of the PPP in a broad range of B cell malignancies that can be efficiently targeted by small molecule inhibition of PP2A and G6PD.

Source URL: <https://www.cirm.ca.gov/about-cirm/publications/b-cell-specific-diversion-glucose-carbon-utilization-reveals-unique>